

Personal Protective Equipment Guidelines

LEVEL I PROTECTION*

- Administrative/Clerical
- Admissions
- Care Management
- Central Supply/Equipment Distribution
- Finance
- Food & Nutrition (cashier; Starbucks)
- IS
- Medical Records
- Non-clinical caregiver (not patient-facing)
- PBX/Call Center
- Pharmacy
- Plant Operations/Maintenance
- Quality/Safety/Infection Prevention/Regulatory

Use Instructions

Fabric/Personal Mask — Home launder before use and as needed and maintain in clean condition.

Hospital Issue Mask — Reuse. Replace if it becomes visibly soiled, wet, contaminated or damaged.

Personal Masks are not allowed in Isolation rooms. Place Isolation mask for appropriate situations.

**Unless you are in direct patient care — within 6 feet of a patient you must use Level 2 protection; discuss with your core leader*

LEVEL II PROTECTION

Direct Patient Care

- Behavioral Health
- Chaplains/Spiritual Care
- Check-in Screeners
- Child Life Specialists
- Clinics
- ECHO/PV
- EEG/EKG
- Emergency Department
- EMS/Ambulance Service
- EVS
- Greeters
- Hydro Therapy/Wound Care
- Infusion Areas
- Lab (draw stations)
- Nursing
- Patient Navigators
- PeriOp Area
- Providers (Physicians/Surgeons/NP/PA, etc.)
- PT/OT/Speech
- Radiology
- Respiratory
- Retail Pharmacy
- Security
- Transportation
- Valet

Use Instructions

» MUST BE WORN UNDER FACE SHIELD «

Hospital-issue Mask — Reuse. Replace if it becomes visibly soiled, wet, contaminated or damaged. *Discard mask in reprocessing box*

Fabric with Filter Pocket Mask — Must be worn with filter. Replace if it becomes visibly soiled, wet, contaminated or damaged. Place mask in bin for professional laundering and dispose of used filter in trash.

Face Shield — Remove shield when no longer in direct contact with patient.

- Disinfect face shield between each use with wipes (No bleach) and place in paper bag with name.
- Use hand hygiene and remove mask.
- Store in clean bag.
- Do not write on mask or shield.

LEVEL III PROTECTION

Procedural

- Cath Lab
- Endoscopy
- Isolation Room (Per policy — example: TB)
- OR

Use Instructions

» MUST BE WORN UNDER FACE SHIELD «

N95 Hospital-issue Mask — *Use in Procedure areas only.* Replace mask if becomes visibly soiled, wet, contaminated or damaged.

Face Shield — Remove shield when no longer in direct contact with patient.

- Disinfect face shield between each use with wipes (No bleach) and place in paper bag with name.
- Use hand hygiene and remove mask.
- Store in clean bag.
- Do not write on mask or shield.

COVID-19 PROTECTION

PAPR/CAPR protection to be utilized on any PUI or confirmed COVID-19 patient for all direct patient care.